

LISTING OF THE CLAIMS

- 1-5. (Canceled).
6. (Previously Presented) An isolated polypeptide comprising:
- (a) the amino acid sequence of the polypeptide of SEQ ID NO: 110;
 - (b) the amino acid sequence of the polypeptide of SEQ ID NO: 110, lacking its associated signal peptide; or
 - (c) the amino acid sequence of the polypeptide encoded by the full-length coding sequence of the cDNA deposited under ATCC accession number 203535.
7. (Previously Presented) The isolated polypeptide of Claim 6 comprising the amino acid sequence of the polypeptide of SEQ ID NO: 110.
8. (Previously Presented) The isolated polypeptide of Claim 6 comprising the amino acid sequence of the polypeptide of SEQ ID NO: 110, lacking its associated signal peptide.
- 9-10. (Canceled).
11. (Original) The isolated polypeptide of Claim 6 comprising the amino acid sequence of the polypeptide encoded by the full-length coding sequence of the cDNA deposited under ATCC accession number 203535.
12. (Previously Presented) A chimeric polypeptide comprising a polypeptide according to Claim 6 fused to a heterologous polypeptide.
13. (Previously Presented) The chimeric polypeptide of Claim 12, wherein said heterologous polypeptide is a tag polypeptide or an Fc region of an immunoglobulin.
14. (Previously Presented) An isolated polypeptide having at least 95% amino acid sequence identity to:
- (a) the amino acid sequence of the polypeptide of SEQ ID NO: 110;
 - (b) the amino acid sequence of the polypeptide of SEQ ID NO: 110, lacking its associated signal peptide; or
 - (c) the amino acid sequence of the polypeptide encoded by the full-length coding sequence of the cDNA deposited under ATCC accession number 203535;
- wherein said isolated polypeptide or a fragment thereof can be used to generate an antibody which can be used to specifically detect the polypeptide of SEQ ID NO: 110 in esophageal tissue samples.

Appl. No. : **10/063,617**
Filed : **May 3, 2002**

15. (Previously Presented) The isolated polypeptide of Claim 14 having at least 99% amino acid sequence identity to:

(a) the amino acid sequence of the polypeptide of SEQ ID NO: 110;

(b) the amino acid sequence of the polypeptide of SEQ ID NO: 110, lacking its associated signal peptide; or

(c) the amino acid sequence of the polypeptide encoded by the full-length coding sequence of the cDNA deposited under ATCC accession number 203535;

wherein said isolated polypeptide or a fragment thereof can be used to generate an antibody which can be used to specifically detect the polypeptide of SEQ ID NO: 110 in esophageal tissue samples.

16. (Previously Presented) A chimeric polypeptide comprising a polypeptide according to Claim 14 fused to a heterologous polypeptide.

17. (Previously Presented) The chimeric polypeptide of Claim 16, wherein said heterologous polypeptide is a tag polypeptide or an Fc region of an immunoglobulin.